## Recombinant Mouse Semaphorin-4C/SEMA4C Protein (Fc Tag)

## Catalog Number: PKSM041138

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description    |  |
|----------------|--|
| Species        | Mouse  |
| Source         | HEK293 Cells-derived Mouse Semaphorin-4C/SEMA4C protein Ala21-Gly664, with an            |
|                | C-terminal Fc  |
| Calculated MW  | 99.3 kDa   |
| Observed MW    | 115-125 kDa  |
| Accession      | Q64151   |
| Bio-activity   | Not validated for activity   |
| Properties     |  |
| Purity         | > 95 % as determined by reducing SDS-PAGE.   |
| Endotoxin      | < 1.0 EU per µg of the protein as determined by the LAL method.                          |
| Storage        | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 |
|                | °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of      |
|                | reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.                      |
| Shipping       | This product is provided as lyophilized powder which is shipped with ice packs.          |
| Formulation    | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.                              |
|                | Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants         |
|                | before lyophilization.   |
|                | Please refer to the specific buffer information in the printed manual.                   |
| Reconstitution | Please refer to the printed manual for detailed information.                             |

Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Semaphorin-4C is a protein which belongs to the semaphorin family, contains 1 Ig-like C2-type domain, 1 PSI domain, 1 Sema domain. As cell surface receptor for PLXNB2, it plays an important role in cell-cell signaling. PLXNB2 binding promotes downstream activation of RHOA and phosphorylation of ERBB2 at 'Tyr-1248'. It required for normal brain development, axon guidance and cell migration, Probable signaling receptor which may play a role in myogenic differentiation through activation of the stress-activated MAPK cascade.