IL-9/Cytokine P40 (C-6His), Human, Recombinant

Cat. No. : PCK206

General Information

Synonyms	Interleukin-9;IL-9;Cytokine P40;T-Cell Growth Factor P40;IL9
Species	Human
Expression host	Human Cells
Sequence	Gln19-Ile144
Accession	P15248
Tag	C-6His
Mol mass	15.2 kDa
Expiration date	12 months

Product feature

Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin (EU/μg)	< 0.1
Storage	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
Shipping	Ice bag
Formulation	Lyophilized from a 0.2 μm filtered solution of 20 mM PB, 150 mM NaCl, pH 7.4.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Background

Interleukin-9 (IL-9) is a secreted Protein that belongs to the IL-7/IL-9 family. IL-9 supports IL-2 independent and IL-4 independent growth of helper T-cells. IL-9 stimulates cell proliferation and prevents apoptosis. It functions through the IL-9 Receptor (IL-9R), which activates different signal transducer and activator (STAT) Proteins and thus connects this Cytokine to various biological processes. IL-9 has been identified as a candidate gene for asthma. IL-9 is a determining factor in the pathogenesis of bronchial hyperresponsiveness.

SDS-PAGE

