

## Recombinant E.coli Tryptophan Synthase $\alpha$ Chain/Trp A Protein

Catalog Number: PKSQ050056

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	E.coli
<b>Source</b>	E.coli-derived E.coli Tryptophan Synthase $\alpha$ Chain/Trp A protein Met1-Ser268
<b>Calculated MW</b>	28.7 kDa
<b>Observed MW</b>	27 kDa
<b>Accession</b>	P0A877
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per $\mu$ g of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 $\mu$ m filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Tryptophan synthase is an enzyme that catalyzes the final two steps in the biosynthesis of tryptophan. It is commonly found in Eubacteria, Archaeobacteria, Protista, Fungi, and Plantae, but is absent from animals such as humans.

Tryptophan synthase typically exists as an  $\alpha$ - $\beta$ - $\alpha$  complex. The alpha subunit is responsible for the aldol cleavage of indoleglycerol phosphate to indole and glyceraldehyde 3-phosphate: L-serine + 1-C-(indol-3-yl)glycerol 3-phosphate = L-tryptophan + D-glyceraldehyde 3-phosphate + H<sub>2</sub>O. The beta subunits catalyze the irreversible condensation of indole and serine to form tryptophan in a pyridoxal phosphate (PLP) dependent reaction. Their assembly into a complex leads to structural changes in both subunits resulting in reciprocal activation.

### For Research Use Only