

## ACOX3 Polyclonal Antibody

catalog number: E-AB-14480

**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*

### Description


|                     |                                                                                    |
|---------------------|------------------------------------------------------------------------------------|
| <b>Reactivity</b>   | Human                                                                              |
| <b>Immunogen</b>    | Recombinant protein of human ACOX3                                                 |
| <b>Host</b>         | Rabbit                                                                             |
| <b>Isotype</b>      | Rabbit IgG                                                                         |
| <b>Purification</b> | Affinity purification                                                              |
| <b>Buffer</b>       | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |

### Applications


### Recommended Dilution

|     |            |
|-----|------------|
| IHC | 1:50-1:200 |
|-----|------------|

### Data



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using ACOX3 Polyclonal Antibody at dilution 1:35



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using ACOX3 Polyclonal Antibody at dilution 1:35

### Preparation & Storage

|                 |                                                                                                          |
|-----------------|----------------------------------------------------------------------------------------------------------|
| <b>Storage</b>  | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.                                          |
| <b>Shipping</b> | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

### Background

Acyl-Coenzyme A oxidase 3 also known as pristanoyl -CoA oxidase (ACOX3) is involved in the desaturation of 2-methyl branched fatty acids in peroxisomes. Unlike the rat homolog, the human gene is expressed in very low amounts in liver such that its mRNA was undetectable by routine Northern-blot analysis or its product by immunoblotting or by enzyme activity measurements. However the human cDNA encoding a 700 amino acid protein with a peroxisomal targeting C-terminal tripeptide S-K-L was isolated and is thought to be expressed under special conditions such as specific developmental stages or in a tissue specific manner in tissues that have not yet been examined.

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.7