Recombinant Rat Fcy RIIB/CD32b Protein(His Tag)

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Catalog Number: PDMR100064



Description Species Rat Source Mammalian-derived Rat Fcy RIIB/CD32b proteins His32-Pro212, with an C-terminal His Mol Mass 19.8 kDa O63203 Accession **Bio-activity** Not validated for activity **Properties** Purity >90% as determined by reducing SDS-PAGE. Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 Storage °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at $< -20^{\circ}$ C for 3 months. Shipping This product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Formulation Mannitol. Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data

	R
j	11
	1 1 1 J 3

SDS-PAGE analysis of Rat Fcγ RIIB/CD32b proteins , 2µg/lane of Recombinant Rat Fcγ RIIB/CD32b proteins was resolved with SDS-PAGE under reducing conditions , showing bands at 40 KD

Background

The Fc gamma Rs have been divided into three classes based on close relationships in their extracellular domains; these groups are designated Fc gamma RI (also known as CD64), Fc gamma RII (CD32), and Fc gamma RIII (CD16). Each group may be encoded by multiple genes and exist in different isoforms depending on species and cell type. The CD64 proteins are high affinity receptors (~10e-8-10e-9 M) capable of binding monomeric IgG, whereas the CD16 and CD32 proteins bind IgG with lower affinities (~10e-6-10e-7 M) only recognizing IgG aggregates surrounding multivalent antigens.

For Research Use Only