

Recombinant Human VEGF-A/VEGF121 Protein (His Tag)

Catalog Number: PKSH031983

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human VEGF-A;VEGF121 protein Ala27-Arg147, with an C-terminal His
Calculated MW	15.1 kDa
Observed MW	16-18 kDa
Accession	P15692-9
Bio-activity	Immobilized Human VEGFR1-Fc(PKSH033445) at 5µg/ml (100 µl/well) can bind Human VEGF 121-His(PKSH031983).The ED ₅₀ of Human VEGF 121-His(PKSH031983)is 9.44 ng/ml.


Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.



Immobilized Human VEGFR1-Fc(PKSH033445) at 5µg/ml (100 µl/well) can bind Human VEGF 121-His(PKSH031983).The ED₅₀ of Human VEGF 121-His(PKSH031983)is 9.44 ng/ml.

Background

For Research Use Only

Human VEGF121; also known as Vascular endothelial growth factor A; VEGFA; Vascular permeability factor; VPF and VEGF; is a homodimeric, heparin-binding glycoprotein which belongs to the platelet-derived growth factor (PDGF)/vascular endothelial growth factor (VEGF) family. VEGF-A is a glycosylated mitogen that specifically acts on endothelial cells and has various effects; including mediating increased vascular permeability; inducing angiogenesis; vasculogenesis; permeabilization of blood vessels and endothelial cell growth; increasing microvascular permeability; promoting cell migration and inhibiting apoptosis. Alternatively spliced transcript variants of VEGF-A encode either secreted or cell-associated isoforms. The lymphangiogenesis may be promoted by upregulation of VEGF121; which may in turn act in part via induction of VEGF-C. It binds to the FLT1/VEGFR1 and KDR/VEGFR2 receptors; heparan sulfate and heparin. NRP1/Neuropilin-1 binds isoforms VEGF-165 and VEGF-145. Isoform VEGF165B binds to KDR but does not activate downstream signaling pathways; does not activate angiogenesis and inhibits tumor growth.

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017