## Recombinant Mouse Igfbp6 Protein(His Tag)

Catalog Number: PDMM100053



Note: Centrifuge before opening to ensure complete recovery of vial contents.

		crip				
	00	CI	РΤ	n	tπ	Λn
JU.	$\mathbf{c}$	v.		w	ш	UП

Species Mouse

**Source** Mammalian-derived Mouse Igfbp6 proteins Ala26-Gly238, with an C-terminal His

 Mol\_Mass
 23.3 kDa

 Accession
 P47880

**Bio-activity** Not validated for activity

## **Properties**

**Purity** > 90% as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping

This product is provided as lyophilized powder which is shipped with ice packs.

Formulation


Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%

Mannitol

**Reconstitution** It is recommended that sterile water be added to the vial to prepare a stock solution of

0.5 mg/mL. Concentration is measured by UV-Vis.

#### Data



SDS-PAGE analysis of Mouse Igfbp6 proteins, 2 µg/lane of Recombinant Mouse Igfbp6 proteins was resolved with an SDS-PAGE under reducing conditions, showing bands at 23.3KD

### Background

#### For Research Use Only

# Recombinant Mouse Igfbp6 Protein(His Tag)

Catalog Number: PDMM100053



The superfamily of insulin-like growth factor (IGF) binding proteins include the six high-affinity IGF binding proteins (IGFBP) and at least four additional low-affinity binding proteins referred to as IGFBP related proteins (IGFBP-rP). All IGFBP superfamily members are cysteine-rich proteins with an conserved cysteine residues, which are clustered in the amino-and carboxy-terminal thirds of the molecule. IGFBPs modulate the biological activities of IGF proteins. Some IGFBPs may also have intrinsic bioactivity that is independent of their ability to bind IGF proteins. Post-translational modifications of IGFBPs, including glycosylation, phosphorylation and proteolysis, have been shown to modify the affinities of the binding proteins to IGF.

Mouse IGFBP-6 cDNA encodes a 238 amino acid (aa) residue precursor protein with an a putative 25 aa residue signal peptide that is processed to generate the 213 aa residue mature protein that is O-glycosylated. Mouse and Human IGFB P-6 share 73% amino acid similarity. Mouse and rat IGFBP-6 share 94% amino acid similarity and the Mouse IGFBP-6 has a 9 amino acid insertion compared to the rat homolog. IGFBP-6 is expressed in ovarian, testicular, muscle, heart and lung tissues in the adult Mouse. IGFBP-6 was not detected in total RNA from a whole Mouse embryo.