

# Recombinant Human S100A9 Protein

Catalog Number: PKSH033380

Note: Centrifuge before opening to ensure complete recovery of vial contents.

## Description

Species Human

Source E.coli-derived Human S100A9 protein Thr 2-Pro114

 Calculated MW
 13.2 kDa

 Observed MW
 14 kDa

 Accession
 P06702

**Bio-activity** Immobilized Recombinant Human S100A9(PKSH033380) at 2μg/ml (100 μl/well) can

bind Anti-Human S100A9 Antibody. The  $\mathrm{ED}_{50}$  of Anti-Human S100A9 Antibody is

 $0.58\mu g/ml$ .

## **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants


before lyophilization.


Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## Data







Immobilized Recombinant Human S100A9(PKSH033380) at  $2\mu g/ml~(100~\mu l/well)$  can bind Anti-Human S100A9 Antibody. The ED50 of Anti-Human S100A9 Antibody is  $0.58\mu g/ml.$ 

# Background

## For Research Use Only

Web:www.elabscience.com

# Elabscience®

## **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

Protein S100-A9 (also MRP14 and calgranulin B)is a calcium- and zinc-binding protein which plays a prominent role in the regulation of inflammatory processes and immune response. It can induce neutrophil chemotaxis, adhesion, can increase the bactericidal activity of neutrophils by promoting phagocytosis via activation of SYK, PI3K/AKT, and ERK1/2 and can induce degranulation of neutrophils by a MAPK-dependent mechanism.

Toll-free: 1-888-852-8623 Web:www.elabscience.com Tel: 1-832-243-6086 Email:techsupport@elabscience.com Fax: 1-832-243-6017