

## FITC Anti-Human CD4 Antibody[SK3]

Catalog Number: E-AB-F1352C

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description

|                         |                                                                                                                                                          |
|-------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|
| Reactivity              | Human;Rhesus;Cynomolgus                                                                                                                                  |
| Host                    | Mouse                                                                                                                                                    |
| Isotype                 | Mouse IgG1, $\kappa$                                                                                                                                     |
| Clone No.               | SK3                                                                                                                                                      |
| Isotype Control         | FITC Mouse IgG1, $\kappa$ Isotype Control[MOPC-21] [Product E-AB-F09792C]                                                                                |
| Conjugation             | FITC                                                                                                                                                     |
| Conjugation Information | FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical filter centered near 530 nm (e.g., a 525/40 nm bandpass filter). |
| Storage Buffer          | Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.                                                                           |


### Applications

### Recommended usage

#### FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

### Data



Human peripheral blood leucocytes are stained with FITC Anti-Human CD4 Antibody (Left). Leucocytes are stained with FITC Mouse IgG1,  $\kappa$  Isotype Control (Right).

### Preparation & Storage

|          |                                                                                                                                                           |
|----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|
| Storage  | Keep as concentrated solution.<br>This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze. |
| Shipping | Ice bag                                                                                                                                                   |

### Antigen Information

|                 |                                                                 |
|-----------------|-----------------------------------------------------------------|
| Alternate Names | T-cell surface antigen T4/Leu-3;T-cell surface glycoprotein CD4 |
| Uniprot ID      | P01730                                                          |
| Gene ID         | 920                                                             |

### For Research Use Only

## Background

CD4, also known as T4, is a 55 kD single-chain type I transmembrane glycoprotein expressed on most thymocytes, a subset of T cells, and monocytes/macrophages. CD4, a member of the Ig superfamily, recognizes antigens associated with MHC class II molecules and participates in cell-cell interactions, thymic differentiation, and signal transduction. CD4 acts as a primary receptor for HIV, binding to HIV gp120. CD4 has also been shown to interact with IL-16.