

A Reliable Research Partner in Life Science and Medicine

Recombinant SerpinA8/Angiotensinogen/AGT Monoclonal Antibody

catalog number: AN300213P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human SerpinA8 protein

Host Rabbit Isotype lgG Clone B131 **Purification** Protein A


Buffer 0.2 µm filtered solution in PBS

Applications Recommended Dilution


1:500-1:2000 **WB** 1:25-1:100 **FCM** 1:20-1:100 ICC/IF

ΙP 4-6 µL/mg of lysate

Data




Flow cytometric analysis of Human AGT expression on HepG2 cells. The cells were stained with purified anti-Human AGT, then a FITC-conjugated second step antibody. with 1% Triton X-100 in PBS, blocked with 10% serum, and The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of intact cells.



Immunofluorescence analysis of Human AGT in Hela or SKBR3 cells. Cells were fixed with 4% PFA, permeabilzed incubated with Rabbit anti-Human AGT Monoclonal Antibody (1:60). Then cells were stained with the Alexa Fluor® 488-conjugated Goat Anti-rabbit IgG secondary antibody (left panel, captured by laser confocal scanning microscope; right panel, captured by fluorescence microscope), countstained with DAPI for nuclear staining (blue). Positive staining was localized to cytoplasm.





Toll-free: 1-888-852-8623 Fax: 1-832-243-6017 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com Rev. V1.2

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Immunoprecipitation analysis using 2 μ L anti-SerpinA8 Monoclonal Antibody and 15 μ l of 50 % Protein G agarose. Western blot was performed from the immunoprecipitate using SerpinA8 Monoclonal Antibody at a dilution of 1:200.

Lane A:0.5 mg HepG2 Whole Cell Lysate

Observed-MW:51 kDa Calculated-MW:51 kDa

Western Blot with SerpinA8 / Angiotensinogen / AGT Monoclonal Antibody at dilution of 1:500. Lane A: HepG2 Whole Cell Lysate, Lysates/proteins at 30 µg per lane.

Observed-MW:51 kDa Calculated-MW:51 kDa

Preparation & Storage

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

Background

Essential component of the renin-angiotensin system (RAS), a potent regulator of blood pressure, body fluid and electrolyte homeostasis. In response to lowered blood pressure, the enzyme renin cleaves angiotensinogen to produce angiotensin-1 (angiotensin 1-10). Angiotensin-1 is a substrate of ACE (angiotensin converting enzyme) that removes a dipeptide to yield the physiologically active peptide angiotensin-2 (angiotensin 1-8). Angiotensin-1 and angiotensin-2 can be further processed to generate angiotensin-3 (angiotensin 2-8), angiotensin-4 (angiotensin 3-8). Angiotensin 1-7 is cleaved from angiotensin-2 by ACE2 or from angiotensin-1 by MME (neprilysin). Angiotensin 1-9 is cleaved from angiotensin-1 by ACE2. Angiotensin-2 acts directly on vascular smooth muscle as a potent vasoconstrictor, affects cardiac contractility and heart rate through its action on the sympathetic nervous system, and alters renal sodium and water absorption through its ability to stimulate the zona glomerulosa cells of the adrenal cortex to synthesize and secrete aldosterone. Angiotensin-3 stimulates aldosterone release. Angiotensin 1-7 is a ligand for the G-protein coupled receptor MAS1 (By similarity). Has vasodilator and antidiuretic effects (By similarity). Has an antithrombotic effect that involves MAS1-mediated release of nitric oxide from platelets.

For Research Use Only