# **Elabscience**®

## **CDK2AP1** Polyclonal Antibody


#### catalog number: E-AB-92415

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description  |  |
|--------------|--|
| Reactivity   | Human  |
| Immunogen    | Recombinant fusion protein of human CDK2AP1  |
| Host         | Rabbit   |
| Is otype     | IgG  |
| Purification | Affinity purification  |
| Buffer       | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |
| Applications | Decommonded Dilution   |

| Applications | Recommended Dilution |
|--------------|----------------------|
| WB           | 1:500-1:2000         |

#### Data



Western blot analysis of extracts of various cell lines using

CDK2AP1 Polyclonal Antibody at 1:3000 dilution.

### Observed-MV:Refer to figures Calculated-MV:9 kDa/12 kDa

| Preparation & Storage |  |
|-----------------------|--|
| Storage               | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.  |
| Shipping              | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

#### Background

The protein encoded by this gene is a cyclin-dependent kinase 2 (CDK2) -associated protein which is thought to negatively regulate CDK2 activity by sequestering monomeric CDK2, and targeting CDK2 for proteolysis. This protein was found to also interact with DNA polymerase alpha/primase and mediate the phosphorylation of the large p180 subunit, which suggests a regulatory role in DNA replication during the S-phase of the cell cycle. This protein also forms a core subunit of the nucleosome remodeling and histone deacetylation (NURD) complex that epigenetically regulates embryonic stem cell differentiation. This gene thus plays a role in both cell-cycle and epigenetic regulation. Alternative splicing results in multiple transcript variants encoding distinct isoforms.