

## Recombinant Human HDHD2 Protein (His Tag)

**Catalog Number:** PKSH032520

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	E.coli-derived Human HDHD2 protein Met 1-Leu259, with an N-terminal His
<b>Calculated MW</b>	30.7 kDa
<b>Observed MW</b>	30 kDa
<b>Accession</b>	Q9H0R4
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 50mM NaCl, pH 8.0. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
<b>Reconstitution</b>	Please refer to the specific buffer information in the printed manual.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Haloacid Dehalogenase-Like Hydrolase Domain-Containing Protein 22 (HDHD2) is a member of the HAD-like hydrolase superfamily. HDHD2 includes L-2-Haloacid Dehalogenase, Epoxide Hydrolases and Phosphatases. There are two active sites in HDHD2 - an L-2-Haloacid Dehalogenase and a Carboxylate group. The L-2-Haloacid Dehalogenase active site catalyzes the hydrolytic dehalogenation of D- and L-2-Haloalkanoic Acids, producing L- and D-2-Hydroxyalkanoic Acids.

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017