

Recombinant Bovine S100A7/PSOR1 Protein (His Tag)

Catalog Number: PDEB100011

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Bovine
Source	E.coli-derived Bovine S100A7 protein Ser2-Gln101, with an N-terminal His
Calculated MW	10.9 kDa
Observed MW	11 kDa
Accession	Q28050
Bio-activity	Not validated for activity

Properties

Purity	> 90% as determined by reducing SDS-PAGE.
Endotoxin	< 10 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Bovine S100A7/PSOR1 proteins, 2 µg/lane of Recombinant Bovine S100A7/PSOR1 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 11 kDa.

Background

S100A7 is a 11-12 kDa member of the S100 family of EF hand calcium binding proteins. Human S100A7 shares 32% amino acid sequence identity with mouse S100A7A, the closest related protein in mouse. It is acetylated at the N-terminus and binds both calcium and zinc ions. S100A7 is up-regulated in keratinocytes of psoriasis and atopic dermatitis lesions, as well as in epithelial cells of the tongue, eye, and female genital tract. Its up-regulation can be induced by bacterial exposure, inflammatory cytokines, or epidermal barrier disruption. S100A7 supports epithelial integrity through killing *E. coli* by sequestration of zinc and through inducing the up-regulation of tight junction proteins. The interaction of S100A7 with RAGE promotes the migration of immune cells and the infiltration of macrophages into tumor sites.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017