

Recombinant Human IDE/Insulysin Protein (His Tag)

Catalog Number: PKSH032593

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human IDE;Insulysin protein Met42-Leu1019, with an C-terminal His
Calculated MW	114.3 kDa
Observed MW	120 kDa
Accession	P14735
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Concentration	Subject to label value.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/ gel packs. Upon receipt, store it immediately at < - 20°C.
Formulation	Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, 0.05% Brij35, 10% Glycerol, pH 7.5.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Insulin-Degrading Enzyme (IDE) is a secreted enzyme that belongs to the peptidase M16 family. IDE is a large zinc-binding protease and cleaves multiple short polypeptides that vary considerably in sequence. IDE plays a role in the cellular breakdown of insulin, IAPP, glucagon, bradykinin, kallidin, and other peptides, and thereby plays a role in intercellular peptide signaling. IDE degrades amyloid formed by APP and IAPP. IDE may participate in the degradation and clearance of naturally secreted amyloid β -protein by neurons and microglia. IDE, which migrates at 110 kDa during gel electrophoresis under denaturing conditions, has since been shown to have additional substrates, including the signaling peptides glucagon, TGF α and β -endorphin.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017