

A Reliable Research Partner in Life Science and Medicine

# **ACADSB Polyclonal Antibody**

catalog number: E-AB-66388

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Human; Mouse; Rat

Immunogen Recombinant fusion protein of human ACADSB

Host Rabbit
Isotype IgG


**Purification** Affinity purification

**Buffer** Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.


## Applications Recommended Dilution

**IHC** 1:50-1:200


#### Data



Immunohistochemistry of paraffin-embedded rat brain using ACADSB Polyclonal Antibody at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining



Immunohistochemistry of paraffin-embedded human breast cancer using ACADSB Polyclonal Antibody at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry of paraffin-embedded mouse spinal cord using ACADSB Polyclonal antibody at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.

#### **Preparation & Storage**

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

**Shipping** The product is shipped with ice pack upon receipt, store it immediately at the

temperature recommended.

### Background

#### For Research Use Only

#### **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

Short/branched chain acyl-CoA dehydrogenase(ACADSB) is a member of the acyl-CoA dehydrogenase family of enzymes that catalyze the dehydrogenation of acyl-CoA derivatives in the metabolism of fatty acids or branch chained amino acids. Substrate specificity is the primary characteristic used to define members of this gene family. The ACADSB gene product has the greatest activity towards the short branched chain acyl-CoA derivative, (S)-2-methylbutyryl-CoA, but also reacts significantly with other 2-methyl branched chain substrates and with short straight chain acyl-CoAs. The cDNA encodes for a mitochondrial precursor protein which is cleaved upon mitochondrial import and predicted to yield a mature peptide of approximately 43.7-KDa.

Fax: 1-832-243-6017