

A Reliable Research Partner in Life Science and Medicine

PE Anti-Mouse CD69 Antibody[H1.2F3]

Catalog Number: E-AB-F1187UD

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse

Host Armenian Hamster
Isotype Armenian Hamster IgG

Clone No. H1.2F3

Isotype Control PE Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F09853D]

Conjugation PE

Conjugation Information PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green

(561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42

nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage


FCM Each lot of this antibody is quality control tested by flow cytometric analysis. Please

check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the

reagent to obtain optimal results [The recommended concentration is 0.1-1 μ g/10⁶ cells in 400 kH column 1

in 100 µL volume].

Data



Cell Stimulation Cocktail-stimulated C57BL/6 mouse splenocytes (5 h) are stained with PE Anti-Mouse CD69 Antibody (filled gray histogram). Unstained stimulated splenocytes (empty black histogram) are used as control.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names AIM;CLEC2C;EA1;GP32/28;MLR-3;VEA

 Uniprot ID
 P37217

 Gene ID
 12515

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Background

CD69 is a 60 kD type II membrane protein composed of a 27/33 kD disulfide-linked homodimer, also known as Very Early Activation Antigen (VEA), AIM, EA1, MLR3, and gp34/28. It is expressed on a subset of thymocytes and platelets. CD69 is rapidly induced on activated T and B cells, neutrophils, and NK cells. It is a C-type lectin, closely related to the NKR-P1 and Ly-49 NK cell activation molecules. CD69 is involved in the early events of cell activation and thymocyte positive selection.

Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Email:techsupport@elabscience.com Web:www.elabscience.com