

A Reliable Research Partner in Life Science and Medicine

Elab Fluor® 488 Anti-Rat CD4(domain 1) Antibody[OX-38]

Catalog Number: E-AB-F1105L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Rat
Host Mouse

Isotype Mouse IgG2a, κ

Clone No. OX-38

Isotype Control Elab Fluor® 488 Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802L]

Conjugation Elab Fluor® 488

Conjugation Information Elab Fluor® 488 is designed to be excited by the Blue laser (488 nm) and detected using

an optical filter centered near 520 nm (e.g., a 525/40 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Rat splenocytes are stained with Elab Fluor[®] 488 Anti-Rat CD4(domain 1) Antibody (filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD4;T-cell surface antigen T4/Leu-3;T-cell surface glycoprotein CD4;domain 1

 Uniprot ID
 P05540

 Gene ID
 24932

For Research Use Only



Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

Background

CD4, also known as T4, is a 55kD glycoprotein member of the immunoglobin superfamily and is expressed on majority of thymocytes, macrophages, and a peripheral T cell subset (T helper cells). CD4 is a T cell co-receptor that interacts with the MHC class II molecule and is involved in T cell activation.