Recombinant Osteonectin/SPARC Monoclonal Antibody

catalog number: AN300525P


Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description

Reactivity	Mouse
Immunogen	Recombinant Mouse Osteonectin/SPARC protein
Host	Rabbit
Isotype	IgG
Clone	B455
Purification	Protein A
Buffer	0.2 μm filtered solution in PBS

Applications	Recommended Dilution
WB	1:500-1:1000
IP	0.2-1 μL/mg of lysate


Data



Western Blot with SPARC Monoclonal Antibody at dilution of 1:500 dilution. Lane A: A549 Whole Cell Lysate, Lysates/proteins at 30 μg per lane.

Observed-MW:40 kDa

Calculated-MW:35 kDa



Immunoprecipitation analysis using 0.5 μL anti-Mouse SPARC Monoclonal Antibody and 15 μl of 50 % Protein G agarose. Western blot was performed from the immunoprecipitate using SPARC Monoclonal Antibody at a dilution of 1:500. Lane A:0.5 mg A549 Whole Cell Lysate, Lane B:0.5 mg 293T Whole Cell Lysate

Observed-MW:40 kDa

Calculated-MW:35 kDa

Preparation & Storage

Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Shipping	Ice bag

Background

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com

Secreted protein acidic and rich in cysteine (SPARC), also known as Osteonectin (ON), is a member of the SPARC family. SPARC consists of three domains: an EF-hand domain, a follistatin-like domain and a Kazal-like domain, and each of which has independent activity and unique properties. The activity of SPARC is context- and cell-type-dependent, which is highlighted by the fact that SPARC has shown seemingly contradictory effects on tumor progression in both clinical correlative studies and in animal models. The location of SPARC in the nuclear matrix of certain proliferating cells, but only in the cytosol of postmitotic neurons, indicates potential functions of SPARC as a nuclear protein, which might be involved in the regulation of cell cycle progression and mitosis. It functions not only to modulate cell-cell and cell-matrix interactions, but its de-adhesive and growth inhibitory properties in non-transformed cells have led to studies to assess its role in cancer. Its divergent actions reflect the complexity of this protein, because in certain types of cancers, such as melanomas and gliomas, SPARC is associated with a highly aggressive tumor phenotype, while in others, mainly ovarian, neuroblastomas and colorectal cancers, SPARC may function as a tumor suppressor. Recent studies have also demonstrated a role for SPARC in sensitizing therapy-resistant cancers. Notably, SPARC is linked to human obesity.

Toll-free: 1-888-852-8623
Web: www.elabscience.com
Tel: 1-832-243-6086
Email: techsupport@elabscience.com
Fax: 1-832-243-6017