

IL-32 Monoclonal Antibody(Detector)

catalog number: AN001830P

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human
Immunogen	Recombinant Human IL-32 protein expressed by E.coli
Host	Rat
Isotype	Rat IgG1
Clone	4D9
Purification	Protein A/G Purification
Buffer	Phosphate buffered solution, pH 7.2, containing 0.05% Proclin300.

Applications Recommended Dilution

ELISA Detector	0.1-0.4 µg/mL
-----------------------	---------------

Data



Sandwich ELISA-Recombinant Human IL-32 protein standard curve. Background subtracted standard curve using IL-32 antibody(AN000680P)(Capture), IL-32 Antibody(AN001830P)(Detector) in sandwich ELISA. The reference range value for Recombinant Human IL-32 protein is 15.63-1000 pg/mL.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

For Research Use Only

IL-32 is a recently discovered cytokine that induces various proinflammatory cytokines (TNF-alpha, IL-1beta, IL-6) and chemokines in both human and mouse cells through the NF-kappaB and p38 MAPK inflammatory signal pathways. It is regulated robustly by other major proinflammatory cytokines and is crucial to inflammation and immune responses. Four of the IL-32 isoforms (alpha, beta, gamma, and delta) are the most representative IL-32 transcripts, and the gamma isoform of IL-32 is the most active, although all isoforms are biologically active. IL-32, a cytokine produced mainly by T, natural killer, and epithelial cells induces significant amounts of TNFalpha and MIP-2 and increases the production of both cytokines in a dose-dependent manner. IL-32 has been implicated in inflammatory disorders, Mycobacterium tuberculosis infections, inflammatory bowel disease, and influenza A virus infection, as well as in some autoimmune diseases, such as rheumatoid arthritis, ulcerative colitis, and in the human stomach cancer, human lung cancer, and breast cancer tissues. Thus, IL-32 expression might be valuable as a biomarker for cancer.