

## Purified Anti-Human CD64 Antibody[10.1]


catalog number: E-AB-F1082A

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description

<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Isotype</b>	Mouse IgG1, $\kappa$
<b>Clone</b>	10.1
<b>Buffer</b>	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

### Data



Human peripheral blood monocytes were stained with 0.2 $\mu$ g Purified Anti-Human CD64 Antibody[10.1] (Right) and 0.2 $\mu$ g mouse IgG1, $\kappa$  Isotype Control (Left), followed by Elab Fluor® 647-conjugated goat Anti-mouse IgG Secondary Antibody, then anti-human CD14 FITC-conjugated Monoclonal Antibody.

### Preparation & Storage

<b>Storage</b>	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
<b>Shipping</b>	Ice bag

### Background

CD64 is a 72 kD single chain type I glycoprotein also known as Fc $\gamma$ RI and FcR I. CD64 is a member of the immunoglobulin superfamily and is expressed on monocytes/macrophages, dendritic cells, and activated granulocytes. The expression can be upregulated by IFN- $\gamma$  stimulation. CD64 binds IgG immune complex. It plays a role in antigen capture, phagocytosis of IgG/antigen complexes, and antibody-dependent cellular cytotoxicity (ADCC).

### For Research Use Only