

Mouse IL-1 β Antibody Pair Set

Catalog No.	E-KAB-0007	Applications	ELISA
Synonyms	IL1B, IL1-BETA, IL1F2, catabolin		

Kit components & Storage

Title	Specifications	Storage
Mouse IL-1 β Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year. Avoid freeze / thaw cycles.
Mouse IL-1 β Detection Antibody (Biotin)	1 vial, 50 μ L	Store at -20°C for one year. Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Product Information

Items		Characteristic (E-KAB-0007)	
		Mouse IL-1 β Capture Antibody	Mouse IL-1 β Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Mouse IL-1 β protein	Recombinant Mouse IL-1 β protein
	Swissprot	P10749	
Product details	Reactivity	Mouse	Mouse
	Host	Rat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50% glycerol, pH 7.4	PBS with 0.04% Proclin 300, 1% protective protein, 50% glycerol, pH 7.4
	Purify	Protein A or G	Antigen Affinity
	Specificity	Detects Mouse IL-1 β in ELISAs.	

For Research Use Only

Applications

Mouse IL-1 β Sandwich ELISA Assay:

	Recommended Concentration/Dilution	Reagent	Images										
ELISA Capture	0.5-4μg/mL	Mouse IL-1β Capture Antibody	 <table><caption>Approximate data points from the standard curve</caption><thead><tr><th>Mouse IL-1β concentration (pg/mL)</th><th>Optical Density</th></tr></thead><tbody><tr><td>10</td><td>0.05</td></tr><tr><td>100</td><td>0.2</td></tr><tr><td>1000</td><td>1.0</td></tr><tr><td>10000</td><td>5.0</td></tr></tbody></table>	Mouse IL-1β concentration (pg/mL)	Optical Density	10	0.05	100	0.2	1000	1.0	10000	5.0
Mouse IL-1β concentration (pg/mL)	Optical Density												
10	0.05												
100	0.2												
1000	1.0												
10000	5.0												
ELISA Detection	1:1000-1:10000	Mouse IL-1β Detection Antibody (Biotin)											

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

The protein encoded by this gene is a member of the interleukin 1 cytokine family. This cytokine is produced by activated macrophages as a proprotein, which is proteolytically processed to its active form by caspase 1 (CASP1/ICE). This cytokine is an important mediator of the inflammatory response, and is involved in a variety of cellular activities, including cell proliferation, differentiation, and apoptosis. The induction of cyclooxygenase-2 (PTGS2/COX2) by this cytokine in the central nervous system (CNS) is found to contribute to inflammatory pain hypersensitivity. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2.

For Research Use Only