

## ZAP70 Polyclonal Antibody

catalog number: **E-AB-52977**

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.



### Description

<b>Reactivity</b>	Human;Mouse
<b>Immunogen</b>	Fusion protein of human ZAP70
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Purification</b>	Antigen affinity purification
<b>Buffer</b>	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

### Applications Recommended Dilution

<b>WB</b>	1:500-1:2000
<b>IHC</b>	1:50-1:300

### Data



Western blot analysis of Jurkat cell Mouse liver tissue HepG2 Immunohistochemistry of paraffin-embedded Human liver cell Mouse kidney tissue lysates using ZAP70 Polyclonal cancer tissue using ZAP70 Polyclonal Antibody at dilution of 1:800 1:60(×200)

**Observed-MV:Refer to figures**

**Calculated-MV:70 kDa**

### Preparation & Storage

<b>Storage</b>	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
<b>Shipping</b>	The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

### Background

Zeta-associated protein (ZAP)-70,a member of the Syk family of tyrosine kinases,plays an important role in T-cell receptor signaling,natural killer (NK) cell activation and early B-cell development. ZAP-70 is expressed in a subset of cases of chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL) with unmutated immunoglobulin heavy-chain variable region (IgVH) genes and is associated with poor clinical outcome.

### For Research Use Only