



A Reliable Research Partner in Life Science and Medicine

Biotin Anti-Mouse CD48 Antibody[HM48-1]

Catalog Number: E-AB-F1017B

Note: Centrifuge before opening to ensure complete recovery of vial contents.

_			- 4		
	es	O PI		$\mathbf{I} \cap$	m
ш,		7 I	II JI	II.U.	

Reactivity Mouse

Host Armenian Hamster
Isotype Armenian Hamster IgG

Clone No. HM48-1

Isotype Control Biotin Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F09853B]

Conjugation Biotin

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. For flow

cytometric staining, the suggested use of this reagent is \leq 1.0 μ g per 10⁶ cells in 100 μ L volume or 100 μ L of whole blood. It is recommended that the reagent be titrated for

optimal performance for each application.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names BCM1 surface antigen;BLAST-1;CD48;CD48 antigen;Cd48;HM48-1;MRC OX-45

surface antigen; SLAMF2; sgp-60

 Uniprot ID
 P18181

 Gene ID
 12506

Background CD48 is a 45 kD GPI-anchored glycoprotein also known as BCM1, Blast-1 (human),

Web: www.elabscience.cn

and OX-45 (rat). It is a member of the lg superfamily, expressed on T and B cells and monocytes/macrophages. It plays a role in adhesion and T cell recognition. The

primary ligands for CD48 are CD2 and CD244.