

## Recombinant Human CD31/PECAM1 Protein (Fc Tag)

**Catalog Number:** PKSH033567

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human CD31/PECAM1 protein Gln28-Lys601, with an C-terminal Fc
<b>Calculated MW</b>	91.6 kDa
<b>Observed MW</b>	110-130 kDa
<b>Accession</b>	AAH22512.1
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Semaphorin-4G is the least characterized of the seven known Class 4 transmembrane semaphorin glycoproteins. Class 4 semaphorins play multiple roles in cell attraction or repulsion, such as development of nerve pathways in the brain, promoting or inhibiting proliferation, in some cases organizing immune cell interactions. Semaphorin-4G can be expressed early in development in the central and peripheral nervous systems and in sensory organs, such as cochlea, olfactory epithelium, vomeronasal organ and retina. In adults, Semaphorin-4G can be found in liver, kidney and brain. The human Semaphorin-4G precursor consists of a 17 amino acids signal sequence, a 658 amino acids extracellular domain, a 21 amino acids transmembrane domain, a 142 amino acids cytoplasmic domain with one Ser/Thr phosphorylation site.

### For Research Use Only