

IL-4Ra/CD124 (C-mFc), Human, Recombinant

Cat. No. : PCK198

General Information

Synonyms	Interleukin-4 Receptor subunit alpha;IL-4 Receptor subunit alpha;IL-4R subunit alpha;IL-4 R-alpha;IL-4RA;CD124;IL-4-binding Protein;IL4-BP;IL4R;IL4RA
Species	Human
Expression host	Human Cells
Sequence	Met26-Gln231
Accession	P24394
Mol mass	50.2 kDa
Expiration date	12 months
Bio activity	Loaded Human IL-4-His on HIS1K Biosensor, can bind Human IL-4RA-mFc with an affinity constant of 0.12 nM as determined in BLI assay.


Product feature

Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin (EU/μg)	< 0.1
Storage	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
Shipping	Ice bag
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


Background

Interleukin 4 Receptor alpha (IL4-Ra) is a widely expressed 140 kDa transmembrane glyco Protein in the class I Cytokine Receptor family. Mature human IL4-Ra consists of a 207 amino acid (aa) extracellular domain (ECD) that contains a Cytokine binding region and one fibronectin type III domain, a 24 aa transmembrane segment, and a 569 aa cytoplasmic domain that contains one Box 1 Motif and one ITIM Motif. IL4-Ra plays an important role in Th2-biased immune responses, alternative macrophage activation, mucosal immunity, allergic inflammation, tumor progression, and atherogenesis. Soluble forms of IL4-Ra, generated by alternate splicing or proteolysis, retain Ligand binding properties and inhibit IL-4 bioactivity.

SDS-PAGE



Bio activity



Loaded Human IL-4-His on HIS1K Biosensor, can bind Human IL-4RA-mFc with an affinity constant of 0.12 nM as determined in BLI assay.

Procell[®]systemProcell[®]system