

## APC Anti-Mouse CD44 Antibody[NIM-R8]

**Catalog Number:** AN00917E

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

Description	
<b>Reactivity</b>	Mouse
<b>Host</b>	Rat
<b>Isotype</b>	Rat IgG2a
<b>Clone No.</b>	NIM-R8
<b>Isotype Control</b>	APC Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832E]
<b>Conjugation</b>	APC
<b>Conjugation Information</b>	APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.
Applications	Recommended usage
<b>FCM</b>	Each lot of this antibody is quality control tested by flow cytometric analysis. <b>The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).</b> Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
Data	
<p>Staining of C57BL/6 murine splenocytes cells with APC Anti-Mouse CD44 Antibody[NIM-R8] (filled gray histogram) or APC Rat IgG2a, κ Isotype Control (empty black histogram). Total viable cells were used for analysis.</p>	
Preparation & Storage	
<b>Storage</b>	<p>Keep as concentrated solution.</p> <p>This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.</p>
<b>Shipping</b>	Ice bag
Antigen Information	
<b>Alternate Names</b>	Hermes;Pgp-1;H-CAM;HUTCH-1;ECMR III;gp85;Ly-24
<b>Uniprot ID</b>	P15379
<b>Gene ID</b>	12505
<b>Background</b>	<p>CD44 is a 80-95 kD glycoprotein also known as Hermes, Pgp1, H-CAM, or HUTCH. It is expressed on all leukocytes, endothelial cells, hepatocytes, and mesenchymal cells. As B and T cells become activated or progress to the memory stage, CD44 expression increases from low or mid levels to high levels. Thus, CD44 has been reported to be a valuable marker for memory cell subsets. High CD44 expression on Treg cells has been associated with potent suppressive function via high production of IL-10. CD44 is an adhesion molecule involved in leukocyte attachment to and rolling on endothelial cells, homing to peripheral lymphoid organs and to the sites of inflammation, and leukocyte aggregation.</p>

### For Research Use Only