



A Reliable Research Partner in Life Science and Medicine

# **Recombinant PEG10 Monoclonal Antibody**

catalog number: AN301959L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### **Description**

Reactivity Human; Rat; Mouse

Immunogen Recombinant human PEG10 fragment

 Host
 Rabbit

 Isotype
 IgG, κ

 Clone
 A675


Purification Protein Apurified

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

## Applications Recommended Dilution

WB 1:500-1:1000

#### Data



Western Blot with PEG10 Monoclonal Antibody at dilution of

1:1000. Lane 1: HepG2, Lane 2: HeLa, Lane 3: Mouse

placenta, Lane 4: Rat placenta Observed-MW:55/100 kDa Calculated-MW:55/100 kDa

## **Preparation & Storage**

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

## **Background**

Paternally expressed gene 10(PEG10) is an imprinted gene which is derived from a Ty3/Gypsy long terminal repeat (LTR) retrotransposon family encoding Gag- and Pol-like domains. Deletion of PEG10 in mice leads to embryonic lethality due to defects in placental formation. PEG10 is aberrantly expressed in several cancer types, including hepatocellular carcinoma, and contributes to tumorigenesis by affecting cell proliferation, apoptosis, and metastasis. The PEG10 gene has two overlapping open reading frames that are regulated by the programmed process of -1 frameshifting. The first encoded protein ORF1 (or RF1) has a Gag domain with coiled-coil domain and zinc finger domains, while ORF1-2 (or RF1/RF2) is produced by -1 frameshifting and creates a fusion of the ORF1 Gag domain with a carboxyl-terminal Pol-like protease domain. PEG10 has retained the ability to form virus-like particles (VLPs) that are secreted as small extracellular vesicles delivered to distant sites.

# For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.0