

## PE/Elab Fluor® 594 Anti-Human IL-12/IL-23 p40 Antibody[C8.6]

Catalog Number: AN00843P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Isotype</b>	Mouse IgG1, κ
<b>Clone No.</b>	C8.6
<b>Isotype Control</b>	PE/Elab Fluor® 594 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792P]
<b>Conjugation</b>	PE/Elab Fluor® 594
<b>Conjugation Information</b>	PE/Elab Fluor® 594 is designed to be excited by the blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 620 nm (e.g., a 610/20 nm bandpass filter).
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

### Applications

### Recommended usage

<b>FCM</b>	Each lot of this antibody is quality control tested by flow cytometric analysis. <b>The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).</b> Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
------------	---

### Data



IFN-γ-primed and then LPS-stimulated (24h) human peripheral blood monocytes intracellular stained with PE/Elab Fluor® 594 Anti-Human IL-12/IL-23 p40 Antibody[C8.6] (filled gray histogram) or PE/Elab Fluor® 594 IgG1, κ Isotype Control (empty black histogram).

### Preparation & Storage

<b>Storage</b>	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
<b>Shipping</b>	Ice bag

### Antigen Information

<b>Alternate Names</b>	Interleukin-12 p40; Interleukin-23 p40; Cytotoxic lymphocyte maturation factor (CLMF); Natural killer cell stimulatory factor (NKSF); CTL maturation factor (TcMF); T-cell stimulating factor (TSF); AN008430
------------------------	---

### For Research Use Only

<b>Uniprot ID</b>	P29460
<b>Gene ID</b>	3593
<b>Background</b>	<p>The C8.6 antibody reacts with human IL-12 p40 subunit of the IL-12 p70 heterodimer and IL-23 p40 subunit of the IL-23 p19/p40, as well as p40 monomer and homodimer. The C8.6 antibody has been reported to strongly inhibit different biological activities of IL-12, (e.g., IFN-gamma induction, mitogenic effects on PHA blasts, and enhancement of NK cell-mediated cytotoxicity). The C8.6 antibody can neutralize the bioactivity of natural or recombinant IL-12.</p>