



A Reliable Research Partner in Life Science and Medicine

Granulin/GRN/Progranulin Monoclonal Antibody

catalog number: AN200186P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human Granulin protein

 Host
 Mouse

 Isotype
 IgG1

 Clone
 A1038


 Purification
 Protein A

Buffer 0.2 µm filtered solution in PBS

Applications Recommended Dilution

ICC/IF 1:20-1:100

Data



Immunofluorescence analysis of Human GRN in Hela cells.

Cells were fixed with 4% PFA, blocked with 10% serum, and incubated with mouse anti-Human GRN monoclonal antibody (1:60). Then cells were stained with the Alexa Fluor® 488-conjugated Goat Anti-mouse IgG secondary antibody (green) and counterstained with DAPI (blue).

Positive staining was localized to cytoplasm.

Preparation & Storage

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

Background

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.1

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Progranulin, also known as acrogranin, PC cell-derived growth factor (PCDGF) and epithelin/granulin precursor, is a ubiquitously expressed, 88 kDa, secreted glycoprotein. Structurally, it belongs to none of the well-established growth factor families. Mouse Progranulin is 589 amino acids (aa) in length and contains a 17 aa signal sequence and a 572 aa mature region that has four potential sites for N-linked glycosylation. It has a highly repetitive organization, containing seven tandem copies of a 55-57 aa consensus motif that contains 12 conserved cysteine residues: VxCx5-6Cx5CCx8CCx6CCxDx2HCCPx4Cx5-6Cx2. Progranulin is secreted in an intact form or undergoes proteolysis leading to the release of multiple peptides made from the seven tandem repeats, the granulins. Mouse Progranulin shares 87% and 75% aa sequence identity with rat and human Progranulin, respectively. Progranulin is involved in the regulation of cellular proliferation, as well as differentiation, development, and pathological processes. It has been isolated as a differentially expressed gene during mesothelial differentiation, macrophage development, development synovium of rheumatoid arthritis and osteoarthritis, sexual differentiation of the brain, and has also been shown to be a mediator of cartilage proliferation plus of wound response and tissue repair. High levels of Progranulin expression have been found to be associated with several human cancers, and are believed to contribute to tumorigenesis in breast cancer, clear cell renal carcinoma, invasive ovarian carcinoma, glioblastoma, adipocyte teratoma, and multiple myeloma.

For Research Use Only