

A Reliable Research Partner in Life Science and Medicine

Recombinant p21 Monoclonal Antibody

catalog number: AN301027L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Rat

Immunogen Recombinant Human p21 protein

HostRabbitIsotypeIgG,κCloneB778PurificationProtein A

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications Recommended Dilution


IHC 1:100-1:200 **WB** 1:1000-1:5000

Data



Western Blot with Recombinant p21 Monoclonal Antibody at dilution of 1:1000 dilution. Lane A: HCT-116 cells.

Observed-MW:18 kDa Calculated-MW:18 kDa



Immunohistochemistry of paraffin-embedded human cervical carcinoma tissue using Recombinant p21 Monoclonal Antibody at dilution of 1:200.

Rev. V1.1

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

Background

This gene encodes a potent cyclin-dependent kinase inhibitor. The encoded protein binds to and inhibits the activity of cyclin-cyclin-dependent kinase2 or -cyclin-dependent kinase4 complexes, and thus functions as a regulator of cell cycle progression at G1. The expression of this gene is tightly controlled by the tumor suppressor protein p53, through which this protein mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. This protein can interact with proliferating cell nuclear antigen, a DNA polymerase accessory factor, and plays a regulatory role in S phase DNA replication and DNA damage repair. This protein was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of cyclin-dependent kinase2, and may be instrumental in the execution of apoptosis following caspase activation. Mice that lack this gene have the ability to regenerate damaged or missing tissue. Multiple alternatively spliced variants have been found for this gene.

For Research Use Only