

A Reliable Research Partner in Life Science and Medicine

Recombinant Phospho-Tau (Ser202, Thr205) Monoclonal Antibody

catalog number: AN300378L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen A synthetic phosphopeptide corresponding to residues around

Host Rabbit
Isotype IgG
Clone B304
Purification Protein A


Buffer 10 mM sodium HEPES, 150 mM NaCl, 100 µg/mL protein protectant, 50% glycerol,

pH 7.5

Applications Recommended Dilution

WB 1:2000-1:10000


Data





Western blot analysis of extracts from mouse brain (line A); treated with antigen-specific phosphopeptide (line B) or antigen-specific peptide (line C) using Phospho-Tau (Ser202, Thr205) rabbit monoclonal Antibody at 1:2000 dilution.. (Validation Experiment)

Observed-MW:50-80 kDa Calculated-MW:78 kDa



Western blot analysis of extracts from rat brain using Phospho-Tau (Ser202, Thr205) rabbit monoclonal Antibody at 1:5000 dilution..

Observed-MW:50-80 kDa Calculated-MW:78 kDa

Rev. V1.1

Western blot analysis of 200 ng Recombinant Human Tau Protein (Full Length) (line A) and 200 ng Recombinant Human Tau Protein (Full Length), GSK3beta-phosphorylated (line B) using Phospho-Tau (Ser202, Thr205) Rabbit Monoclonal Antibody at 1:5000 dilution..

> Observed-MW:50-80 kDa Calculated-MW:78 kDa

Preparation & Storage

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com



Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping lce bag

Background

This gene encodes the microtubule-associated protein tau (MAPT) whose transcript undergoes complex, regulated alternative splicing, giving rise to several mRNA species. MAPT transcripts are differentially expressed in the nervous system, depending on stage of neuronal maturation and neuron type. MAPT gene mutations have been associated with several neurodegenerative disorders such as Alzheimer's disease, Pick's disease, frontotemporal dementia, cortico-basal degeneration and progressive supranuclear palsy.

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.1