

MCTS1 Polyclonal Antibody

catalog number: E-AB-15193

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity	Human
Immunogen	Recombinant protein of human MCTS1
Host	Rabbit
Isotype	Rabbit IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications

Recommended Dilution


WB	1:200-1:1000
IHC	1:100-1:300

Data




Western Blot analysis of Jurkat and PC3 cell using MCTS1 Polyclonal Antibody at dilution of 1:200

Calculated-MW:21 kDa



Immunohistochemistry of paraffin-embedded Human brain using MCTS1 Polyclonal Antibody at dilution of 1:60



Immunohistochemistry of paraffin-embedded Human thyroid cancer using MCTS1 Polyclonal Antibody at dilution of 1:60

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.8

MCTS1 (malignant T cell amplified sequence 1), also known as MCT1, is a 181 amino acid protein that is ubiquitously expressed and localizes to the cytoplasm of cells. MCTS1 may play a role in cell cycle regulation by decreasing cell doubling time and by shortening the duration of G1 transit time and G1/S transition. The protein level of Cdk4 and Cdk6 kinases activity and cyclin D1 is enhanced by expression of MCTS1. As a translation enhancer, MCTS1 recruits DENR and binds to the cap complex of the 5'-terminus of mRNAs, subsequently altering the mRNA translation profile. MCTS1 promotes lymphoid tumor development and contributes to the pathogenesis and progression of breast cancer. MCTS1 positively regulates phosphorylation of ERK1 and ERK2.