

PUS10 Polyclonal Antibody

catalog number: E-AB-13544

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description



Reactivity	Human;Mouse
Immunogen	Synthetic peptide of human PUS10
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications

Recommended Dilution

WB	1:200-1:1000
IHC	1:50-1:200


Data



Western Blot analysis of Mouse intestinum tenue tissue using PUS10 Polyclonal Antibody at dilution of 1:200

Immunohistochemistry of paraffin-embedded Human breast cancer using PUS10 Polyclonal Antibody at dilution of 1:45

Calculated-MV:60 kDa



Immunohistochemistry of paraffin-embedded Human brain using PUS10 Polyclonal Antibody at dilution of 1:45

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

Pseudouridination, the isomerization of uridine to pseudouridine, is the most common posttranscriptional nucleotide modification found in RNA and is essential for biologic functions such as spliceosome biogenesis. Pseudouridylate synthases, such as PUS10, catalyze pseudouridination of structural RNAs, including transfer, ribosomal, and splicing RNAs. These enzymes also act as RNA chaperones, facilitating the correct folding and assembly of tRNAs.

For Research Use Only