


Applications

Porcine GDF15 Sandwich ELISA Assay

	Recommended Concentration/Dilution	Reagent	Images																		
ELISA Capture	0.5-4 µg/mL	Porcine GDF15 Capture Antibody	 <p>The graph displays a standard curve for the Porcine GDF15 Sandwich ELISA Assay. The x-axis represents Porcine GDF15 Concentration in pg/mL on a logarithmic scale from 1 to 10,000. The y-axis represents Optical Density on a logarithmic scale from 0.01 to 10. The data points show a clear upward trend, indicating that as the concentration of Porcine GDF15 increases, the optical density also increases proportionally.</p> <table border="1"> <caption>Approximate data points from the standard curve</caption> <thead> <tr> <th>Porcine GDF15 Concentration (pg/mL)</th> <th>Optical Density</th> </tr> </thead> <tbody> <tr> <td>10</td> <td>0.05</td> </tr> <tr> <td>20</td> <td>0.1</td> </tr> <tr> <td>50</td> <td>0.2</td> </tr> <tr> <td>100</td> <td>0.3</td> </tr> <tr> <td>200</td> <td>0.5</td> </tr> <tr> <td>500</td> <td>1.0</td> </tr> <tr> <td>1000</td> <td>2.0</td> </tr> <tr> <td>2000</td> <td>3.0</td> </tr> </tbody> </table>	Porcine GDF15 Concentration (pg/mL)	Optical Density	10	0.05	20	0.1	50	0.2	100	0.3	200	0.5	500	1.0	1000	2.0	2000	3.0
Porcine GDF15 Concentration (pg/mL)	Optical Density																				
10	0.05																				
20	0.1																				
50	0.2																				
100	0.3																				
200	0.5																				
500	1.0																				
1000	2.0																				
2000	3.0																				
ELISA Detection	1:1000-1:10000	Porcine GDF15 Detection Antibody (Biotin)																			

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Bone morphogenetic proteins (e.g. , BMP9 , MIM 605120) are members of the transforming growth factor-beta (see TGFB1 , MIM 190180) superfamily and regulate tissue differentiation and maintenance. They are synthesized as precursor molecules that are processed at a dibasic cleavage site to release C-terminal domains containing a characteristic motif of 7 conserved cysteines in the mature protein. GDF15 mRNA is most abundant in the liver , with lower levels seen in some other tissues. Its expression in liver can be significantly up-regulated in during injury of organs such as liver , kidney , heart and lung.

For Research Use Only