

PSMA3 Polyclonal Antibody

catalog number: E-AB-52718

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human;Mouse;Rat
Immunogen	Fusion protein of human PSMA3
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications

Recommended Dilution

WB	1:500-1:2000
IHC	1:40-1:200


Data




Western blot analysis of NIH/3T3 A549 HL60 and PC3 cell
Mouse liver tissue Mouse spleen tissue Hela cell lysates
using PSMA3 Polyclonal Antibody at dilution of 1:200

Observed-MV:Refer to figures

Calculated-MV:28 kDa



Immunohistochemistry of paraffin-embedded Human thyroid
cancer tissue using PSMA3 Polyclonal Antibody at dilution
of 1:30(×200)



Immunohistochemistry of paraffin-embedded Human
ovarian cancer tissue using PSMA3 Polyclonal Antibody at
dilution of 1:30(×200)

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

Background

For Research Use Only

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. Two alternative transcripts encoding different isoforms have been identified.