A Reliable Research Partner in Life Science and Medicine

# **ADORA1 Polyclonal Antibody**

catalog number: E-AB-92078

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Human; Mouse; Rat

Immunogen A synthetic peptide of human ADORA1

Host Rabbit
Isotype IgG


**Purification** Affinity purification

**Buffer** Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

## **Applications** Recommended Dilution


WB 1:500-1:2000 IHC 1:50-1:200

## Data




Western blot analysis of extracts of Mouse heart using ADORA1 Polyclonal Antibody at 1:1000 dilution.

Observed-MV:37 kDa Calculated-MV:13 kDa/36 kDa



Immunohistochemistry of paraffin-embedded rat brain using ADORA1 Polyclonal Antibody at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry of paraffin-embedded human esophageal using ADORA1 Polyclonal Antibody at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.

## Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

## Background

## For Research Use Only

Toll-free: 1-888-852-8623 Web:www.elabscience.com

Tel: 1-832-243-6086 Email:techsupport@elabscience.com

## **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

The protein encoded by this gene is an adenosine receptor that belongs to the G-protein coupled receptor 1 family. There are 3 types of adenosine receptors, each with a specific pattern of ligand binding and tissue distribution, and together they regulate a diverse set of physiologic functions. The type A1 receptors inhibit adenylyl cyclase, and play a role in the fertilization process. Animal studies also suggest a role for A1 receptors in kidney function and ethanol intoxication. Transcript variants with alternative splicing in the 5' UTR have been found for this gene.

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u>

Tel: 1-832-243-6086 Email:techsupport@elabscience.com