

Recombinant Human LRAP/ERAP2 Protein (His Tag)

Catalog Number: PKSH031454

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human LRAP/ERAP2 protein Ala 56-Thr 960, with an N-terminal His
Calculated MW	106 kDa
Observed MW	115-125 kDa
Accession	NP_071745.1
Bio-activity	Measured by its ability to cleave the fluorogenic peptide substrate, Arg-7-amido-4-methylcoumarin (Arg-AMC). The specific activity is > 50 pmoles/min/μg.

Properties

Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 12.5mM Tris, 75mM NaCl, pH 7.5, 50% glycerol Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Leukocyte-derived arginine aminopeptidase (LRAP), also known as endoplasmic reticulum-aminopeptidase 2 (ERAP2), is the second identified aminopeptidase localized in the in the luminal side of endoplasmic reticulum (ER) processing antigenic peptides presented to major histocompatibility complex (MHC) class I molecules. It is a 960-amino acid protein with significant homology to placental leucine aminopeptidase and adipocyte-derived leucine aminopeptidase. LRAP preferentially hydrolyzes the basic residues Arg and Lys, and contains the HEXXH(X)18E zinc-binding motif, which is the characteristic of the M1 family of zinc metallopeptidases which also includes PILS/ARTS1/ERAP1 and LNPEP/PLAP. Induced by interferon-gamma, LRAP is able to trim various MHC class I antigenic peptide precursors.