Purified Anti-Mouse CD38 Antibody[NIMR5]

catalog number: E-AB-F1193A



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| - | | | | | |
|--------------|----|-----|---|----|----|
| | AC | cri | m | Т | ٦n |
| \mathbf{L} | | CII | | 2. | |


Reactivity Mouse **Host** Rat

buffer PBS, pH 7.2. Contains 0.05% proclin 300.

Applications Recommended Dilution

FCM $2 \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$

Data



C57/BL6 Mouse splenocytes were stained with 0.2 µg Purified Anti-Mouse CD38 Antibody[NIMR5] (Right) and 0.2 µg Rat IgG2a, κ Isotype Control (Left), followed by FITC-conjugated Goat Anti-Rat IgG Secondary Antibody.

Preparation & Storage

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping Ice bag

Background

CD38 is a 42 kD glycoprotein, also known as T10. It is an ADP-ribosyl hydrolase, expressed on B cells, NK cells, a subset of T cells, brain, muscle, and kidney. In mouse, CD38 expression is downregulated on germinal center B cells and plasma cells, whereas this is not the case for humans. By functioning as both a cyclase and a hydrolase, CD38 mediates lymphocyte activation, as well as adhesion and metabolism of cADPR and NAADP. CD31 is the ligand of CD38.

For Research Use Only