

## Recombinant Human VSIG8 Protein (Fc Tag)

Catalog Number: PKSH033221

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human VSIG8 protein Val22-Gly263, with an C-terminal Fc
<b>Calculated MW</b>	54.2 kDa
<b>Observed MW</b>	57 kDa
<b>Accession</b>	P0DPA2
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
<b>Reconstitution</b>	Please refer to the specific buffer information in the printed manual.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

V-set and immunoglobulin domain-containing protein 8 (VSIG8) is a single-pass type I membrane protein. The human VSIG8 cDNA encodes 414 amino acids (aa) including a 21 aa signal sequence; a 242 aa extracellular domain (ECD) containing 2 Ig-like V-type (immunoglobulin-like) domains; a 21 aa transmembrane domain and a 130 aa cytoplasmic domain. The function of VSIG8 is not clear.

### For Research Use Only