

## Recombinant Human MECP2 Protein (His Tag)

**Catalog Number:** PKSH032751

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human MECP2 protein Met 1-Ser486, with an C-terminal His
<b>Calculated MW</b>	53.5 kDa
<b>Observed MW</b>	90 kDa
<b>Accession</b>	P51608
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 90 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM Histidine-HCl, 8% Sucrose, 50mM NaCl, 0.02% Tween 80, pH 6.0. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 90 % as determined by reducing SDS-PAGE.

### Background

The MeCP2 helps regulate gene activity (expression) by modifying chromatin, the complex of DNA and protein that packages DNA into chromosomes. The MeCP2 protein is present in cells throughout the body, although it is particularly abundant in brain cells. In the brain, the MeCP2 protein likely plays a role in maintaining connections (synapses) between neurons, where cell-to-cell communication occurs. The alternative splicing of proteins is critical for normal communication between neurons and may also be necessary for the function of other types of brain cells.

### For Research Use Only