

# MMP-8 Monoclonal Antibody(Capture)

catalog number: AN002080P

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## Description


<b>Reactivity</b>	Human
<b>Immunogen</b>	Recombinant Human MMP-8 protein expressed by Mammalian
<b>Host</b>	Rat
<b>Isotype</b>	Rat IgG1
<b>Clone</b>	2F4
<b>Purification</b>	Protein A/G Purification
<b>Conjugation</b>	Unconjugated
<b>buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.05% proclin 300.

## Applications

## Recommended Dilution

<b>ELISA Capture</b>	2-8 µg/mL
----------------------	-----------

## Data



Sandwich ELISA-Recombinant Human MMP-8 protein standard curve. Background subtracted standard curve using MMP-8 antibody(AN002080P)(Capture), MMP-8 antibody(AN002090P)(Detector) in sandwich ELISA. The reference range value for Recombinant Human MMP-8 protein is 0.16-10 ng/mL.

## Preparation & Storage

<b>Storage</b>	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
<b>Shipping</b>	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

## Background

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMPs are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. However, the enzyme encoded by this gene is stored in secondary granules within neutrophils and is activated by autolytic cleavage. Its function is degradation of type I, II and III collagens. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3.

## For Research Use Only