

Purified Anti-Human CD11b Antibody[ICRF44]

catalog number: E-AB-F11460P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


| | |
|---------------------|--|
| Reactivity | Human |
| Immunogen | Recombinant Human CD11b protein |
| Host | Mouse |
| Isotype | Mouse IgG1, κ |
| Clone | ICRF44 |
| Purification | >98%, Protein A/G purified |
| Conjugation | Unconjugated |
| Buffer | Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling. |

Applications

Recommended Dilution

| | |
|------------|---|
| FCM | 2 $\mu\text{g/mL}$ (1×10^5 - 5×10^5 cells) |
|------------|---|

Data



Human peripheral blood monocytes were stained with 0.2 μg Purified Anti-Human CD11b Antibody[ICRF44] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by APC-conjugated Goat Anti-Mouse IgG Secondary Antibody, then anti-Human CD14 FITC-conjugated Monoclonal Antibody.

Preparation & Storage

| | |
|-----------------|--|
| Storage | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. |
| Shipping | Ice bag |

Background

Human ITGAM (Integrin alpha M subunit) is a 127kDa (predicted) glycoprotein, a member of the Integrin family of proteins. The Integrin family proteins are heterodimeric transmembrane receptors composed of an alpha and a beta subunit. The Integrin alpha M subunit, also known as MAC-1 alpha subunit or CD11b, combines with the Integrin beta 2 subunit (CD18) to form the non-covalent heterodimer Integrin alpha M/ beta 2, also known as MAC-1 and complement receptor type 3 (CR3).

For Research Use Only