Elabscience®

Monkey GDF15 Antibody Pair Set

Catalog No.E-KAB-0669ApplicationsELISASynonymsGDF-15;MIC-1;MIC1;NAG-1;PDF;PLAB;PTGFB;TGF-PL

Kit components & Storage

Title	Specifications	Storage
Monkey GDF15 Capture Antibody	1 vial, 100 µ g	Store at -20° C for one year.
		Avoid freeze/thaw cycles.
Monkey GDF15 Detection Antibody	1 vial, 50 μL	Store at -20° C for one year.
(Biotin)		Avoid freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0669)	
		Monkey GDF15 Capture Antibody	Monkey GDF15 Detection Antibody (Biotin)
Immunogen	Immunogen	Recombinant Monkey GDF15	Recombinant Monkey GDF15 protein
Information		protein	
	Swissprot	G7PWZ3	
Product details	Reactivity	Monkey	Monkey
	Host	Goat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%
		glycerol; pH 7.5	protective protein; 50% glycerol; pH
			7.5
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Monkey GDF15 in ELISAs.	

For Research Use Only

Elabscience®

Applications

Monkey GDF15 Sandwich ELISA Assay:

	Recommended Concentration/Dilution	Reagent	Images
ELISA	0.5-4 μg/mL	Monkey GDF15 Capture	
Capture		Antibody	10
ELISA	1:1000-1:10000	Monkey GDF15 Detection	Optical Density
Detection		Antibody (Biotin)	Ö 0.1
Dettetion			•

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Bone morphogenetic proteins (e.g., BMP9, MIM 605120) are members of the transforming growth factor-beta (see TGFB1, MIM 190180) superfamily and regulate tissue differentiation and maintenance. They are synthesized as precursor molecules that are processed at a dibasic cleavage site to release C-terminal domains containing a characteristic motif of 7 conserved cysteines in the mature protein. GDF15 mRNA is most abundant in the liver, with lower levels seen in some other tissues. Its expression in liver can be significantly up-regulated in during injury of organs such as liver, kidney, heart and lung.