

A Reliable Research Partner in Life Science and Medicine

HDAC5 Polyclonal Antibody

catalog number: E-AB-63255

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat

Immunogen A synthetic peptide of human HDAC5 (NP 005465.2).

Host Rabbit
Isotype IgG


Purification Affinity purification


Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications Recommended Dilution


IHC 1:50-1:200 **IF** 1:50-1:200


Data





Immunohistochemistry of paraffin-embedded Rat liver using Immunohistochemistry of paraffin-embedded Human tonsil HDAC5 Polyclonal Antibody at dilution of 1:100 (40x lens). using HDAC5 Polyclonal Antibody at dilution of 1:100 (40x lens).





Immunohistochemistry of paraffin-embedded Mouse spinal cord using HDAC5 Polyclonal Antibody at dilution of 1:100 (40x lens).

Immunofluorescence analysis of A-549 cells using HDAC5 Polyclonal Antibody

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u>

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to the class II histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. It coimmunoprecipitates only with HDAC3 family member and might form multicomplex proteins. It also interacts with myocyte enhancer factor-2 (MEF2) proteins, resulting in repression of MEF2-dependent genes. This gene is thought to be associated with colon cancer. Two transcript variants encoding different isoforms have been found for this gene.

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u>

Tel: 1-832-243-6086 Email:techsupport@elabscience.com Fax: 1-832-243-6017