

A Reliable Research Partner in Life Science and Medicine

## AF/LE Purified Anti-Mouse CD38 Antibody[NIMR5]

catalog number: E-AB-F11930

Note: Centrifuge before opening to ensure complete recovery of vial contents.

$\mathbf{r}$			
	PC	cri	ption
$\mathbf{L}$	CO	CII	JULUI

**Reactivity** Mouse **Host** Rat


Buffer Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method

.

Applications	Recommended Dilution
	1te commende de la commencia d

FCM  $2 \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$ 

## Data



C57/BL6 Mouse splenocytes were stained with 0.2  $\mu$ g AF/LE

Purified Anti-Mouse CD38 Antibody[NIMR5] (Right) and 0.2  $\mu$ g Rat IgG2a,  $\kappa$  Isotype Control (Left), followed by

FITC-conjugated Goat Anti-Rat IgG Secondary Antibody.

## Preparation & Storage

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles. This preparation contains no preservatives, thus it should be handled

under aseptic conditions.

**Shipping** Ice bag

## Background

CD38 is a 42 kD glycoprotein, also known as T10. It is an ADP-ribosyl hydrolase, expressed on B cells, NK cells, a subset of T cells, brain, muscle, and kidney. In mouse, CD38 expression is downregulated on germinal center B cells and plasma cells, whereas this is not the case for humans. By functioning as both a cyclase and a hydrolase, CD38 mediates lymphocyte activation, as well as adhesion and metabolism of cADPR and NAADP. CD31 is the ligand of CD38.