

## Coagulation Factor III/Tissue Factor/CD142 Monoclonal Antibody

catalog number: AN200028P

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


|                     |                                                      |
|---------------------|------------------------------------------------------|
| <b>Reactivity</b>   | Human                                                |
| <b>Immunogen</b>    | Recombinant Human F3 / Tissue factor / CD142 protein |
| <b>Host</b>         | Mouse                                                |
| <b>Isotype</b>      | Mouse IgG2a                                          |
| <b>Clone</b>        | A872                                                 |
| <b>Purification</b> | Protein A                                            |
| <b>Buffer</b>       | 0.2 µm filtered solution in PBS                      |

### Applications


### Recommended Dilution

|               |            |
|---------------|------------|
| <b>ICC/IF</b> | 1:20-1:100 |
| <b>FCM</b>    | 1:25-1:100 |

### Data



Flow cytometric analysis of Human F3(CD142) expression on A431 cells. Cells were stained with purified anti-Human F3(CD142), then a FITC-conjugated second step antibody. The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of intact cells.



Immunofluorescence analysis of Human F3 in A431 cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with mouse anti-Human F3 Monoclonal Antibody (1:60) at 4°C overnight. Then cells were stained with the Alexa Fluor® 488-conjugated Goat Anti-mouse IgG secondary antibody(green) and counterstained with DAPI for nuclear staining(blue). Positive staining was localized to cell membrane and cytoplasm.

### Preparation & Storage

|                 |                                                                                                                                                                                                                                                    |
|-----------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Storage</b>  | This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles. |
| <b>Shipping</b> | Ice bag                                                                                                                                                                                                                                            |

### Background

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.3

This gene encodes coagulation factor III which is a cell surface glycoprotein. This factor enables cells to initiate the blood coagulation cascades, and it functions as the high-affinity receptor for the coagulation factor VII. The resulting complex provides a catalytic event that is responsible for initiation of the coagulation protease cascades by specific limited proteolysis. Unlike the other cofactors of these protease cascades, which circulate as nonfunctional precursors, this factor is a potent initiator that is fully functional when expressed on cell surfaces. There are 3 distinct domains of this factor: extracellular, transmembrane, and cytoplasmic. This protein is the only one in the coagulation pathway for which a congenital deficiency has not been described. Alternate splicing results in multiple transcript variants.