# **USP12 Polyclonal Antibody**

catalog number: E-AB-18513



Note: Centrifuge before opening to ensure complete recovery of vial contents.

## Description

**Reactivity** Human; Mouse

**Immunogen** Full length fusion protein

Host Rabbit
Isotype IgG


**Purification** Antigen affinity purification

**Conjugation** Unconjugated

**buffer** Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications	Recommended Dilution	
WB	1:500-1:2000	
IHC	1:25-1:100	

#### Data



Western blot analysis of Mouse liver tissue lysate using USP12 Polyclonal Antibody at dilution of 1:700

Immunohistochemistry of paraffin-embedded Human tonsil tissue using USP12 Polyclonal Antibody at dilution of 1:35(×200)

Observed-MV:Refer to figures
Calculated-MV:43 kDa

### Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

### Background

The ubiquitin (Ub) pathway involves three sequential enzymatic steps that facilitate the conjugation of Ub and Ub-like molecules to specific protein substrates. Through the use of a wide range of enzymes that can add or remove ubiquitin, the Ub pathway controls many intracellular processes such as signal transduction, transcriptional activation and cell cycle progression. USP12 (ubiquitin specific peptidase 12), also known as UBH1 or USP12L1, is a 370 amino acid protein belonging to the peptidase C19 family and the USP12/USP46 subfamily. Considered a deubiquitinating enzyme, it is suggested that USP12 has almost no deubiquitinating activity by itself and requires the interaction with WDR48 to have high activity. The gene encoding USP12 maps to human chromosome 13, which houses over 400 genes, such as BRCA2 and RB1, and comprises nearly 4% of the human genome.

### For Research Use Only