Recombinant Human LAMP2/CD107b Protein (His Tag)

Catalog Number: PKSH032684



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-					
- 1	00	cri	m	17	٦m
J			.,,	, T. U	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,

 Species
 Human

 Mol_Mass
 39.4 kDa

 Accession
 P13473

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Lysosomal Associated Membrane Protein 2 (LAMP2) is a major component of lysosomal membranes. LAMP2 is a transmembrane glycoprotein about 110kDa. Mature human LAMP2 consists of a 347 amino acid (aa) intralumenal domain, a 24 aa transmembrane segment, and a 35 aa cytoplasmic tail. The lumenal domain is organized into two heavily N-glycosylated regions. Alternate splicing generates a human LAMP2 isoform (LAMP2B) with a substituted juxtamembrane lumenal region, cytoplasmic tail and transmenmbrane segment.LAMP2 itself can cleavage lysosomal luminal domain and degradation lysosomal. In the help of chaperone HSC73,LAMP2 mediates the lysosomal uptake in complex with cargo proteins and is required for the lysosomal destruction of autophagic vacuoles.

For Research Use Only