Purified Anti-Human CD1a Antibody[Ab571]

catalog number: E-AB-F13730P




Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--------------|--|
| Reactivity | Human |
| Immunogen | Recombinant Human CD1a protein |
| Host | Mouse |
| Isotype | Mouse IgGl, κ |
| Clone | Ab571 |
| Purification | >98%, Protein A/G purified |
| Conjugation | Unconjugated |
| buffer | PBS, pH 7.2. Contains 0.05% proclin 300. |
| Applications | Recommended Dilution |

| FCM | $2 \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$ |
|-----|---|

Data



Molt-4 were stained with 0.2 μ g Purified Anti-Human CD1a

Antibody[Ab571] (Right) and 0.2 μg Mouse IgG1, κ Isotype

Control (Left), followed by FITC-conjugated Goat Anti-

Mouse IgG Secondary Antibody.

| Preparation & Storage | |
|-----------------------|---|
| Storage | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / |
| | thaw cycles. |
| Shipping | Ice bag |
| | |

Background

CD1a is a 49 kDa transmembrane glycoprotein in the CD1 family of glycolipid antigen-presenting MHC-like molecules. CD1a contains one Ig-like domain in its extracellular region. It is expressed by most nonhuman mammals but not by mice or rats. Complexes of CD1a with beta 2-microglobulin and endogenous glycolipids are constitutively expressed on antigen presenting cells, cortical thymocytes, and Langerhans cells. CD1a is a target of autoreactive Th22 helper T cells in the skin.

For Research Use Only