



A Reliable Research Partner in Life Science and Medicine

## **Recombinant Dectin-1/CLEC7A Monoclonal Antibody**

catalog number: AN300495P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Mouse

Immunogen Recombinant Mouse Dectin-1/CLEC7A protein


Host Rabbit
Isotype IgG
Clone B425
Purification Protein A

Buffer 0.2 µm filtered solution in PBS

Applications Recommended Dilution

ICC/IF 1:20-1:100

#### Data



Immunofluorescence analysis of Mouse Dectin-1 in Raw264.7 cells. Cells were fixed with 4% PFA, blocked with 10% serum, and incubated with rabbit anti-mouse Dectin-1 monoclonal antibody (1:60) at 37°C 1 hour. Then cells were stained with the Alexa Fluor® 488-conjugated Goat Antirabbit IgG secondary antibody (green) and counterstained with DAPI (blue).

#### **Preparation & Storage**

**Storage** This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

**Background** 

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.1

# Elabscience®

### Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Dectin-1 was recently identified as the most important receptor for beta-glucan. It is a type II transmembrane protein which binds beta-1,3 and beta-1,6 glucans, and is expressed on most cells of the innate immune system and has been implicated in phagocytosis as well as killing of fungi by macrophages, neutrophils and dendritic cells. Recognition of beta-glucan by dectin-1 triggers effective immune response, including phagocytosis and proinflammatory factor production, to eliminate infecting fungi, which especially benefits immunocompromised patients against opportunistic fungal infection. In addition, dectin-1 is involved in the adaptive immune response as well as autoimmune diseases and immune tolerance. Dectin-1 can recognize and respond to live fungal pathogens and is being increasingly appreciated as having a key role in the innate responses to these pathogens. In addition to its exogenous ligands, Dectin-1 can recognize an unidentified endogenous ligand on T cells and may act as a costimulatory molecule. Recent studies have highlighted the importance of Dectin-1 in anti-fungal immunity, in both mice and humans, and have suggested a possible involvement of this receptor in the control of mycobacterial infections.

For Research Use Only